Double dimensional reduction of M5-brane action in Sen formalism

Anajak Phonchantuek¹ and Pichet Vanichchapongjaroen^{1*}

¹The Institute for Fundamental Study "The Tah Poe Academia Institute", Naresuan University, Phitsanulok, 65000, Thailand.

*Corresponding author(s). E-mail(s): pichetv@nu.ac.th; Contributing authors: anajakp61@nu.ac.th;

Abstract

In this paper, we present double dimensional reduction of the complete M5-brane action in the Sen formalism of self-dual form. Although in this formalism the gravity couples to the independent pseudo-2-form and pseudo-3-form fields in a non-standard and very complicated way, the double dimensional reduction on the complete action can be carried out. This is by dualising some components of the pseudo-2-form field and integrating out the pseudo-3-form field. We show explicitly that the double dimensional reduction on a circle gives rise to the complete D4-brane or the complete dual D4-brane action depending on what components of the pseudo-2-form field are dualised. The duality between the D4-brane and dual D4-brane actions are realised in the viewpoint of the Sen formalism as the swapping of the roles of the components of the pseudo-2-form field between those which are dualised and those which remain. We also make a brief outline on how to generalise this to the cases of dimensional reduction on some other spaces as well as how to understand the duality of reduced action from the viewpoint of the Sen formalism. An explicit example for dimensional reduction of the quadratic six-dimensional action on a torus is given and the realisation of its S-duality is also discussed.

1 Introduction

The constructions of the low-energy effective action describing an M5-brane coupled to the background eleven-dimensional supergravity have been known to be non-trivial. This is largely due to the difficulty in the construction of a six-dimensional action for a chiral 2-form field, which is one of the fields in the field content of the M5-brane. The self-duality of its 3-form field strength H makes the straightforward attempt impossible, since $H \wedge *H = 0$ when imposing H = *H in six dimensions. Therefore, the idea is to first work out the six-dimensional action for a chiral 2-form field then generalise to

the complete M5-brane action coupled to back-ground eleven-dimensional supergravity. Various non-trivial approaches which allow this are for example by giving up manifest diffeomorphism invariance [1], [2], [3], [4], [5], or by introducing auxiliary fields which have no dynamics as in the PST formalism [6], [7], [8], [9], [10], [11], [12], [13]. Once the action of the chiral field is constructed, it is usually relatively simpler to extend to the complete M5-brane action. It is also worth mentioning a recent approach [14], [15], [16] in which further auxiliary fields are introduced to lift the limitation on the auxiliary field in the chiral 2-form action in the PST formalism.

Another recent approach is given in [17], [18] which is motivated by string field theory. This approach, which is called the Sen formalism, in fact applies to self-dual fields in 4p + 2 dimensions. The case relevant to us is therefore p=1. In this case, the Sen formalism gives a new perspective on the 2-form field with linear self-dual field strength that the 2-form field is in fact not an independent field. In this approach, the independent fields are a 2-form field P and a 3-form field Q. Both of these fields are non-standard, for example, they transform in an unusual way under diffeomorphism transformation. Furthermore, the field Q is linear self-dual with respect to flat six-dimensional metric even if the six-dimensional spacetime is curved. It is also required that a certain combination of Q and the curved metric gives rise to a 3-form field H which is off-shell self-dual with respect to the curved metric and is closed on-shell. So in the spacetime with trivial topology, H is exact on-shell. Therefore, in the Sen formalism, instead of describing the 2-form field with linear self-dual field strength, one describes an exact linear self-dual 3-form field which is in fact not an independent field.

Studies of various aspects as well as extensions of the Sen formalism are given for example in [19], [20], [21], [22], [23], [24]. Of particular interest to us is the extension [23] to a complete M5-brane action coupled to background eleven-dimensional supergravity. The construction is within the Green-Schwarz formalism in which only the target space has manifest supersymmetry. The constructed action has all the required symmetries for example gauge symmetries, diffeomorphism, and κ -symmetry.

Although the M5-brane action in the Sen formalism has been constructed, further studies are still required in order to better understand the action. One of the important aspects is that the independent fields P and Q are expected to couple to gravity in a non-standard and complicated way since these fields transform in an unusual way under diffeomorphism and the action is non-linear. In order to work towards this goal, one may study dimensional reduction.

It turns out that even the dimensional reduction of the original quadratic action in Sen formalism is readily a challenge. Although it is possible to carry out dimensional reduction on various spaces, the studies [18], [19] are mostly being restricted to the cases in which the uncompact space is flat. So when extending to the M5-brane action, the situation is expected to be even more complicated. Nevertheless, this is exactly the task we focus on in this work.

The aim of this paper is to present double dimensional reduction of the complete M5-brane action in Sen formalism. In particular, we are going to show explicitly that double dimensional reduction on a circle gives rise to the complete D4-brane action or the complete dual D4-brane action, depending on the processes. It turns out that these two processes are related simply by swapping the components of the psuedo-form P. In fact, the swapping is shown to be related to a duality transformation. We also outline how to generalise this to some other spaces provided that the six-dimensional worldvolume is a Cartesian product and that a chosen set of projection operators on 3-form fields satisfy several simple properties.

This paper is organised as follows. In section 2, we review the M5-brane action in the Sen formalism. This action is a complete action of M5-brane in the eleven-dimensional supergravity background. Basic properties of this action are reviewed. In particular, we give a brief explanation on how the coupling of this action to gravity is complicated. In section 3, we present double dimensional reduction of the M5-brane action in the Sen formalism on a circle. We first review and slightly modify the approach of [20] for the dimensional reduction of the Sen quadratic action on a circle. By a further modification, the approach can be applied to the M5-brane action in the Sen formalism whose induced metric is in a general form. The approach involves integrating out some components of the pseudo-form P, while the remaining components are combined with some other fields, through appropriate field redefinitions, to become standard fields. Depending on the way that the components are chosen, one arrives to either the complete D4-brane or the complete dual D4-brane actions each of which is coupled to type IIA supergravity background. The duality between the D4-brane and the dual D4-brane actions can be realised from the viewpoint of the Sen formalism as coming from swapping of the roles of the components of P. With the insight gained from the analysis, we proceed in section 4 to generalise to the cases of dimensional reduction on other spaces. The generalisation looks promising as it gives the reduced action whose physical part describes standard fields. Dualities for reduced actions can also be given from the viewpoint of the Sen formalism. Further checks and generalisations could also be possible which we leave as future works. We give conclusions and discussions in section 5.

2 The M5-brane action with self-dual 3-form

In this section, we review the form of the M5-brane action in the Sen formalism [17], [18] which makes use of non-standard self-dual 3-forms in the sense that the self-duality is with respect to flat metric although the theory may couple to the curved metric.

In the Sen formalism of self-dual (2p+1)-form in 4p+2 dimensional spacetime, there are two kinds of Hodge star operators. The first kind is the standard Hodge star operator * which is defined in the standard way, dependent on the spacetime metric. The second kind is denoted *' which is similar to the standard Hodge star operator but is defined with respect to 4p+2 dimensional Minkowski metric.

From now on, let us focus on the case p = 1 which corresponds to self-dual 3-form in six dimensional spacetime. In this case, the action takes the form [17], [18], [19], [20]

$$S = \int \left(\frac{1}{4}dP \wedge *'dP - Q \wedge dP + \mathcal{L}_I(Q, g, J)\right),$$
(2.1)

where P is a 2-form field, Q is a 3-form field, g is the six dimensional metric, and J is a 3-form external source. Note that we have made the rescaling of the fields with respect to those originally given in [17], [18] in order for the action to have the standard scaling. This is closely related to the scaling adopted in [19]. The field Q is *'-self-dual, that is Q = *'Q. The variation of the action of \mathcal{L}_I with respect to Q takes the form

$$\delta_Q \mathcal{L}_I = \delta Q \wedge R(Q, g, J). \tag{2.2}$$

Therefore, R is *'-anti-self-dual. Define

$$H^J \equiv Q - R + J,\tag{2.3}$$

which should be *-self-dual off-shell. That is, for linear theory, $H^J = *H^J$. The equations of motion for P and Q imply that

$$dH^{(s)} = 0, dH^J = dJ, (2.4)$$

where

$$H^{(s)} = Q + \frac{1}{2}(dP + *'dP). \tag{2.5}$$

The kinetic term of the field P in the action (2.1) has the wrong sign. So the field P is unphysical. However, this is not harmful as the combination $H^{(s)}$ decouples from the physical field H^J . From eq.(2.4), it can be seen that in spacetime with trivial topology, $H^J - J$ is exact on-shell.

It turns out that when promoting to nonlinear theory [23], the only change in the above discussion is simply that the linear *-self-duality becomes the non-linear *-self-duality condition

$$*H^J = \mathcal{V}(H^J, g). \tag{2.6}$$

In particular, a complete M5-brane action in the Green-Schwarz formalism in the form of eq.(2.1) reads [23]

$$S_{M5} = \frac{1}{2} \int \left(\frac{1}{2} dP \wedge *' dP - 2Q \wedge dP - \frac{1}{12} *U(F, g) + Q \wedge R + 2C_6 + F \wedge C_3 \right),$$
(2.7)

which describes a six-dimensional worldvolume embedded in the eleven-dimensional background target superspace. For definiteness, let us call the action (2.7) as the Sen M5-brane action. The fields g, C_3 , and C_6 are induced from the background target superspace. In the context of the action (2.1), the field C_3 is identified as J whereas the field F is identified as H^J . That is

$$F = Q - R + C_3. (2.8)$$

Note that we have rescaled the action eq.(2.7) with respect to that presented in [23]. This is in order for the scaling to better match with other M5-brane actions presented in the literature. The scaling we made with respect to [23] is $C_6 \rightarrow C_6/2, S_{M5} \rightarrow S_{M5}/2$. The explicit form of \mathcal{V} in eq.(2.6) in the context of M5-brane and the form of U_{M5} in the action (2.7) can be described in index

notation. So let us define the convention before coming back to express V and U.

Let middle Greek alphabets for example μ, ν, ρ represent indices of the coordinates of the six-dimensional worldvolume. The worldvolume coordinates are then labelled as dx^{μ} . Define d^6x through

$$dx^{\mu_1} \wedge dx^{\mu_2} \wedge \dots \wedge dx^{\mu_6} = d^6 x \epsilon^{\mu_1 \mu_2 \dots \mu_6}, \quad (2.9)$$

where $\epsilon^{\mu_1\mu_2\cdots\mu_6}$ is the Levi-Civita symbol defined such that $\epsilon^{012345} = 1$. The * operator is defined as

$$*(dx^{\mu_{1}} \wedge \cdots \wedge dx^{\mu_{p}})$$

$$= dx^{\mu_{p+1}} \wedge \cdots \wedge dx^{\mu_{6}} \frac{(-1)^{p+1}}{(6-p)! \sqrt{-g}} \times \qquad (2.10)$$

$$g_{\mu_{p+1}\nu_{p+1}} \cdots g_{\mu_{6}\nu_{6}} \epsilon^{\nu_{p+1}\cdots\nu_{6}\mu_{1}\cdots\mu_{p}}.$$

Similarly, the *' operator is defined as

$$*'(dx^{\mu_1} \wedge \cdots \wedge dx^{\mu_p})$$

$$= dx^{\mu_{p+1}} \wedge \cdots \wedge dx^{\mu_6} \frac{(-1)^{p+1}}{(6-p)!} \times$$

$$\eta_{\mu_{p+1}\nu_{p+1}} \cdots \eta_{\mu_6\nu_6} \epsilon^{\nu_{p+1}\cdots\nu_6\mu_1\cdots\mu_p}.$$

$$(2.11)$$

Differential p-forms are expressed in coordinates as

$$\omega_p = \frac{1}{p!} dx^{\mu_1} \wedge dx^{\mu_2} \wedge \dots \wedge dx^{\mu_p} \omega_{\mu_p \dots \mu_2 \mu_1}. \quad (2.12)$$

Exterior derivative is defined to act from the right

$$d\omega_p = \frac{1}{p!} dx^{\mu_1} \wedge dx^{\mu_2} \wedge \dots \wedge dx^{\mu_p} \wedge dx^{\nu} \partial_{\nu} \omega_{\mu_p \dots \mu_2 \mu_1}.$$
(2.13)

For M5-brane, the non-linear *-self-duality condition (2.6) can be expressed as

$$(*F)_{\mu\nu\rho} = \left(-\frac{U}{12} + \frac{24}{U}\right) F_{\mu\nu\rho} + \frac{6}{U} (F^3)_{\mu\nu\rho},$$
(2.14)

where

$$(F^3)_{\mu\nu\rho} \equiv F_{[\mu|\nu'\rho'}F^{\mu'\nu'\rho'}F_{\mu'|\nu\rho]},$$
 (2.15)

and U, which also appears in the action (2.7), is given by

$$U = -24\sqrt{1 + \frac{F_{\mu\nu\rho}F^{\mu\nu\rho}}{24}}.$$
 (2.16)

The indices in eq.(2.14)-(2.16) are raised by $g^{\mu\nu}$.

The action (2.7) possesses all the required symmetries for example diffeomorphism, gauge symmetries, and kappa-symmetry. It is interesting to note that for each symmetry, transformation rules for P and Q are related by

$$\delta Q = -\frac{1}{2}(1 + \epsilon')d\delta P, \qquad (2.17)$$

which implies that $H^{(s)}$ is invariant under all of the symmetry transformations. This is desirable since $H^{(s)}$ decouples from physical fields. Its symmetry transformations should be zero in order to not affect the symmetry transformations of physical fields. Focusing on diffeomorphism, the transformations on P and Q under $x^{\mu} \mapsto x^{\mu} + \xi^{\mu}$ are

$$\delta_{\xi}P = i_{\xi}(F - C_3), \quad \delta_{\xi}Q = -\frac{1}{2}(1 + *)di_{\xi}(F - C_3),$$
(2.18)

which are non-standard for 2-form and 3-form fields. Due to this, the fields P and Q are called pseudo-forms.

Although the Sen M5-brane action (2.7) is shown to be a complete M5-brane action, further studies are still required in order to better understand this action. Most notably, since P and Q transform in a non-standard way under diffeomorphism, their coupling to gravity are non-standard and is in fact complicated. This complication is readily seen in the quadratic action,

$$S = \int \left(\frac{1}{4}dP \wedge *'dP - Q \wedge dP + \frac{1}{2}Q \wedge \mathcal{M}Q\right),\tag{2.19}$$

which for definiteness, we will call this as the Sen quadratic action. Here, \mathcal{M} is a linear operator on 3-form and is given by 1

$$\mathcal{M} = (1 + **')^{-1}(1 - **'). \tag{2.20}$$

Part of non-standard coupling to gravity is due to the operator **'. Its appearance through $\mathcal M$ makes this non-standard coupling even more complicated since $\mathcal M$ can in principle be expressed as a power series in **' up to the 19th order. Indeed, the complication is even further amplified in the

 $^{^{1}}$ In principle, there are many possible ways to define the operator \mathcal{M} . See [17], [18], [20] for alternative definitions.

Sen M5-brane action as it is non-linear in Q. It is not even clear whether it can be given in the closed form when expressed explicitly in Q.

One way to better understand how the Sen M5-brane action is coupled to gravity is to study dimensional reduction, which provides a simpler scenario to study when compared with the Sen M5-brane action itself.

In [18], [20], dimensional reductions of the Sen quadratic action are carried out on various spaces. Since the action couples to gravity in a complicated way, in each case a special metric is chosen such that the uncompactified space is flat. This makes it possible to carry out the analysis and is sufficient for the purpose of extracting some physics. It was remained to see whether the cases of more general metric can be studied without much difficulties.

In the next section, we will present the study of the double dimensional reduction on a circle of the complete Sen M5-brane action which is coupled to the background eleven-dimensional target superspace. We will demonstrate how to successfully study the full case in which the induced metric takes a general form dependent on the coordinates of reduced worldvolume.

3 Double dimensional reduction on a circle

3.1 Dimensional reduction of the Sen quadratic action

Let us review the analysis of dimensional reduction of the Sen quadratic action (2.19) on a circle. In [20], the analysis is given both in the Hamiltonian and Lagrangian formalisms. Let us however only review the analysis of [20] in the Lagrangian formalism and only follow the key ideas. The exact analysis will be carried out slightly differently especially at the last stage. This results in the interpretation being slightly improved.

Consider the case where the six-dimensional spacetime has the metric of the form

$$g = \begin{pmatrix} \eta_5 & 0\\ 0 & r^2 \end{pmatrix}, \tag{3.1}$$

where η_5 is the five-dimensional flat metric, and r is a positive real number. The coordinate x^5 has length dimension and is compactified as $x^5 \sim$

 $x^5 + l$. Let every field be independent from the coordinate x^5 .

Let Roman alphabets for example a,b,c,i,j,k,m,n represent indices of the coordinates of the five-dimensional spacetime. Denote

$$\epsilon_{ijkmn} \equiv \epsilon_{ijkmn5}, \qquad \epsilon^{ijkmn} \equiv \epsilon^{ijkmn5}.$$
 (3.2)

Therefore, ϵ_{ijkmn} and ϵ^{ijkmn} are Levi-Civita symbols on the five-dimensional spacetime.

Due to *'-self-duality of Q and *'-anti-self-duality of R, each of them has only ten independent components. Let us denote independent components for each of Q and R as

$$q_{ab} \equiv Q_{ab5}, \qquad R_{ab} \equiv R_{ab5}. \tag{3.3}$$

Therefore,

$$Q_{abc} = \frac{1}{2} \epsilon_{abcij} \eta^{im} \eta^{jn} q_{mn},$$

$$R_{abc} = -\frac{1}{2} \epsilon_{abcij} \eta^{im} \eta^{jn} R_{mn}.$$
(3.4)

Let us express R_{ab} in terms of q_{ab} . This can be done by computing

$$R = \mathcal{M}Q,\tag{3.5}$$

which gives

$$R_{ab} = -\frac{r-1}{r+1}q_{ab}. (3.6)$$

The action (2.19) then becomes

$$S = \frac{l}{2} \int d^5x \left(\frac{3}{4} \partial_a P_{bc} \partial^{[a} P^{bc]} + \partial_a P_{b5} \partial^{[a} P^{b]}_5 + 2q^{ab} \partial_a P_{b5} + \frac{1}{2} q_{ab} \partial_m P_{np} \epsilon^{mnpab} + \frac{(r-1)}{(r+1)} q^{ab} q_{ab} \right).$$

$$(3.7)$$

Next, let us make a replacement

$$X_{abc} = 3\partial_{[a}P_{bc]} \tag{3.8}$$

on eq.(3.7) and introduce a Lagrange multiplier K_m which imposes $\partial_{[a}X_{bcd]}=0$. The action then

becomes

$$S = \frac{l}{2} \int d^{5}x \left(\frac{1}{12} X_{abc} X^{abc} + \frac{1}{6} q_{ab} \epsilon^{abijk} X_{ijk} \right.$$

$$+ \frac{1}{6} X_{abc} \partial_{m} K_{n} \epsilon^{abcmn} + \partial_{a} P_{b5} \partial^{[a} P^{b]}_{5}$$

$$+ 2q^{ab} \partial_{a} P_{b5} + \frac{(r-1)}{(r+1)} q^{ab} q_{ab} \right).$$

$$(3.9)$$

By integrating out X followed by q, we obtain

$$S = \frac{l}{2} \int d^5x \left(\frac{r-1}{2r} \partial_a K_b \partial^{[a} K^{b]} - \frac{r+1}{r} \partial_a K_b \partial^{[a} P^{b]5} + \frac{r-1}{2r} \partial_a P_{b5} \partial^{[a} P^{b]}_5 \right).$$
(3.10)

We may view the quantity inside the bracket as a quadratic form whose matrix

$$\frac{1}{2r} \begin{pmatrix} r-1 & -(r+1) \\ -(r+1) & r-1 \end{pmatrix}$$
 (3.11)

has eigenvalues -1/r, 1 and the corresponding eigenvectors $(1,1)^T$, $(1,-1)^T$. Therefore, let us define

$$f_{ab} = l(\partial_{[a}K_{b]} + \partial_{[a}P_{b]}), \quad f_{ab}^{(s)} = l(\partial_{[a}K_{b]} - \partial_{[a}P_{b]})$$

where the factor l is introduced so that f and $f^{(s)}$ have mass dimension two. This corresponds to the field redefinition

$$A_a = \frac{l}{2}(K_a + P_a), \quad A_a^{(s)} = \frac{l}{2}(K_a - P_a).$$
 (3.13)

So f_{ab} and $f_{ab}^{(s)}$ are field strengths of A_a and $A_a^{(s)}$, respectively. With the field redefinitions, the action becomes

$$S = \int d^5x \left(-\frac{1}{4rl} f_{ab} f^{ab} + \frac{1}{4l} f_{ab}^{(s)} (f^{(s)})^{ab} \right),$$
(3.14)

which describes two uncoupled Maxwell fields with opposite signs of kinetic terms. For any value of $r \in \mathbb{R}^+$, the field A_a has the correct sign of the kinetic term with the 1/r scaling as required by

conformal symmetry [25]. On the other hand, the field $A_a^{(s)}$ has the wrong sign of the kinetic term.

Note that the difference between the analysis in [20] and ours is just only in the field redefinition which we have made in eq.(3.13). The counterpart presented in [20] leads to the conclusion that the kinetic term of the standard Maxwell field scales with 1/r only in the limit $r \to 0$. On the other hand, the field redefinition (3.13) leads to the conclusion that the kinetic term of the standard Maxwell field scales with 1/r for any value of $r \in \mathbb{R}^+$, not only in the $r \to 0$ limit. The 1/r scaling upon dimensional reduction on a circle is also in agreement with the result from the PST formalism. As a simple check, consider the quadratic version of the M5-brane action in dual formulation. By considering the six-dimensional spacetime with metric (3.1), identifying x^5 as a coordinate on the circle, and keeping only the zero Fourier mode, the action presented in eq.(3.19) of the reference [5] reduces to the five-dimensional Maxwell theory with the scaling 1/r. The exact agreement with the requirement from conformal symmetry and the relevant result from the PST formalism further confirms that the Sen quadratic action is a quadratic action for an M5-brane.

In the next subsections, we will proceed to extend the idea of this subsection to analyse double dimensional reductions of the Sen M5-brane action on a circle. It will be seen that although the Sen M5-brane action couples to gravity in a complicated way, the analysis based on slight modifications of the steps given in this subsection turns out to apply also for the Sen M5-brane action.

3.2 The complete D4-brane action from the Sen M5-brane action

In the previous subsection, we have reviewed the dimensional reduction of the Sen quadratic action to give a five-dimensional Maxwell theory. We worked with the case where the metric is given by eq.(3.1), in which the six-dimensional worldvolume is a cartesian product of a flat five-dimensional spacetime with a circle with coordinate x^5 .

It is possible, after slightly modifying the strategy, to extend the analysis to the case of the Sen M5-brane action which is a non-linear action with source terms. In the previous subsection, we have seen that the dimensional reduction is obtained

by first expressing R in terms of Q, dualising some components of P, then using Euler-Lagrange equations to eliminate Q. Then after field redefinition, we are left with the action which describes an unphysical field decoupled with Maxwell field in five dimensions.

In this section, since we are considering a non-linear action with source terms, expressing R directly in terms of Q would be too complicated if at all possible. As to be given in details below, it turns out that the modification of the strategy is simply to switch some steps. This would allow us to successfully dimensionally reduce the Sen M5-brane action to the complete action which describes a D4-brane in type IIA supergravity background uncoupled with an unphysical field.

Let us consider a double dimensional reduction of the Sen M5-brane action (2.7) by letting the background target space be compactifized such that X^{10} is on a circle and let the x^5 coordinate of M5-brane worldvolume wraps around X^{10} . The length of x^5 is given by

$$\int dx^5 = 2\pi g_s l_s. \tag{3.15}$$

So if the tension T_{M5} of the M5-brane action (2.7) is introduced back, which is simply by making a scaling $S_{M5} \mapsto T_{M5}S_{M5}$, then $T_{M5}2\pi g_s l_s = T_{D4}$ as expected. So from now on, let us set for simplicity

$$\int dx^5 = 1, \qquad T_{M5} = 1 = T_{D4}. \tag{3.16}$$

Under the double dimensional reduction, the induced metric on the worldvolume is given by

$$g_{\mu\nu} = \begin{pmatrix} e^{-2\phi/3} \gamma_{ab} + e^{4\phi/3} C_a C_b & e^{4\phi/3} C_a \\ e^{4\phi/3} C_b & e^{4\phi/3} \end{pmatrix}.$$
(3.17)

The fields ϕ and C_a are considered as pullbacks of the background superfield to five-dimensional worldvolume. Let us next consider how the pullbacks C_3 and C_6 of the corresponding target space superfields are dimensionally reduced. The components of C_3 are separated via

$$C_3 = \frac{1}{2} dx^a \wedge dx^b \wedge dx^5 C_{ba5} + \frac{1}{3!} dx^i \wedge dx^j \wedge dx^k C_{kij}.$$
(3.18)

It is natural to denote the following:

$$b_{ab} \equiv C_{ab5}, \qquad \tilde{C}^{ab} \equiv \frac{1}{3!} \epsilon^{abijk} C_{ijk}.$$
 (3.19)

Note by definition that \tilde{C}^{ab} is the dualisation of C_{ijk} with respect to five-dimensional flat metric η . So it would be convenient to define raising and lowering of the indices of B and \tilde{C} by using five-dimensional flat metric. As for C_6 , it can be expressed as

$$C_6 = -C_5 \wedge dx^5. \tag{3.20}$$

So in summary, for the reduced theory, the pull-backs of the target space superfields of type IIA supergravity to the five-dimensional worldvolume theory are γ_{ab} , b_2 , C_1 , C_3 , C_5 .

The worldvolume action then reads

$$S = \frac{1}{2} \int d^{5}x \left(\frac{3}{4} \partial_{a} P_{bc} \partial^{[a} P^{bc]} + \partial_{a} P_{b5} \partial^{[a} P^{b]}_{5} \right)$$

$$+ 2q^{ab} \partial_{a} P_{b5} + \frac{1}{2} q_{ab} \partial_{m} P_{np} \epsilon^{mnpab}$$

$$+ \frac{1}{12} \sqrt{-g} U + [q(R+b)]$$

$$- \frac{1}{2} [(q-R)(b-\tilde{C})] - \int C_{5}.$$
(3.21)

Since, unlike the case discussed in the previous subsection, the five-dimensional part of the world-volume is not flat. So one has to be more careful when defining trace as there are two kinds of metric being used here to raise and lower indices. Let us revisit the definition of trace. From now on, it will be understood that components of matrices appearing within the trace are of the form $(\cdot)^a{}_b$. For example,

$$[MN] = M^a{}_b N^b{}_a. (3.22)$$

For a given five-dimensional matrix M, we will specify whether its components are raised by five-dimensional flat metric η or five-dimensional curved metric γ . The components of matrices q, R, B, \tilde{C} which we encounter so far are raised and lowered by η . We will also later define other five-dimensional matrices and will specify the metric which are used to raise or lower their components.

If we were to follow the same strategy as that of the previous section, we would have to eliminate Rfrom eq.(3.21) by expressing it in terms of Q, g, C_3 . Here, however, let us instead keep R explicit at this stage and simply proceed to dualise P_{ab} . By defining X_{abc} as in eq.(3.8) and introducing the Lagrange multiplier term, we obtain

$$S = \frac{1}{2} \int d^{5}x \left(\frac{1}{12} X_{abc} X^{abc} + \partial_{a} P_{b5} \partial^{[a} P^{b]}_{5} + 2q^{ab} \partial_{a} P_{b5} + \frac{1}{6} q_{ab} X_{mnp} \epsilon^{mnpab} + \frac{1}{3!} X_{abc} \partial_{m} K_{n} \epsilon^{abcmn} + \frac{1}{12} \sqrt{-g} U + [q(R+b)] - \frac{1}{2} [(q-R)(b-\tilde{C})] \right) - \int C_{5}.$$
(3.23)

Integrating out X gives

$$S = \frac{1}{2} \int d^5x \left(-\left[(\partial K + \partial P + q)^2 \right] + 2 \left[\partial K \partial P \right] + \frac{1}{12} \sqrt{-g} U + \left[q(R+b) \right] - \frac{1}{2} \left[(q-R)(b-\tilde{C}) \right] \right) - \int C_5,$$
(3.24)

where ∂K and ∂P are matrices with components $(\partial K)_{mn} \equiv \partial_{[m}K_{n]}$ and $(\partial P)_{mn} \equiv \partial_{[m}P_{n]5}$, respectively. The components of ∂K and ∂P are raised and lowered by η . Let us define

$$\psi \equiv \partial K + \partial P, \qquad \psi^{(s)} \equiv \partial K - \partial P, \qquad (3.25)$$

whose components are also raised and lowered by η . We may view the above definition as the field redefinitions

$$K_m + P_m \equiv -\frac{A_m}{2}, \qquad K_m - P_m \equiv \frac{A_m^{(s)}}{2}.$$
 (3.26)

So $-\psi$ and $\psi^{(s)}$ are field strengths of A and $A^{(s)}$. The action becomes

$$S = \frac{1}{2} \int d^5x \left(-\left[(\psi + q)^2 \right] + \frac{1}{2} [\psi^2] - \frac{1}{2} [(\psi^{(s)})^2] + \frac{1}{12} \sqrt{-g} U + [q(R+b)] - \frac{1}{2} [(q-R)(b-\tilde{C})] \right) - \int C_5.$$
(3.27)

Euler-Lagrange equation for q gives

$$R = \psi + q. \tag{3.28}$$

By using eq.(3.28) to eliminate R from the action, we obtain

$$S = \int d^{5}x \left(-\frac{1}{4} [(\psi^{(s)})^{2}] - \frac{1}{4} [\psi^{2}] + \frac{1}{2} [q(-\psi + b)] + \frac{1}{24} \sqrt{-g} U + \frac{1}{4} [\psi b] - \frac{1}{4} [\psi \tilde{C}] \right) - \int C_{5}.$$
(3.29)

It is possible to simplify the expression for U. Let us define matrices F and \tilde{F} whose components are

$$F^{a}_{b} \equiv e^{-2\phi/3} F^{a}_{b5}, \quad \tilde{F}^{a}_{b} \equiv e^{-2\phi/3} \tilde{F}^{a}_{b5}. \quad (3.30)$$

It will be understood that indices of F_{ab} and \tilde{F}_{ab} are raised and lowered by γ . With this convention, we have

$$F_{\mu\nu\rho}F^{\mu\nu\rho} = -3[F^2] + 3[\tilde{F}^2].$$
 (3.31)

Since twenty components of $F_{\mu\nu\rho}$ are related to each other by the non-linear self-duality condition, only ten independent components are required. For our case, we need to eliminate \tilde{F}_{ab5} . The relevant non-linear self-duality condition is

$$\tilde{F}_{ab} = \frac{(1 - y_1)F_{ab} + (F^3)_{ab}}{\sqrt{1 - y_1 + y_1^2 - y_2}},$$
(3.32)

where

$$y_1 \equiv \frac{1}{2}[F^2], \qquad y_2 \equiv \frac{1}{4}[F^4].$$
 (3.33)

Therefore, after substituting into eq.(3.31) and then working out U, we obtain

$$U = -12 \frac{2 - y_1}{\sqrt{1 - y_1 + \frac{1}{2}y_1^2 - y_2}}.$$
 (3.34)

Let us eliminate q. By considering the ijk component of H = Q - R along with eq.(3.28), we obtain

$$q^{ab} = \frac{1}{2} \left(\sqrt{-g} \tilde{F}^{ab} + \tilde{C}^{ab} - \frac{1}{2} \epsilon^{abijk} F_{ij} C_k + \eta^{ac} \eta^{bd} F_{cd} - b^{ab} \right).$$

$$(3.35)$$

This completely determine Q in terms of other fields due to eq.(3.32) and

$$F_{ab} = -\psi_{ab} + b_{ab}, (3.36)$$

where, as defined in eq.(3.25)-(3.26), ψ_{ab} is a field strength of a 1-form field.

So by substituting eq.(3.35) into eq.(3.29), the action becomes

$$S = \int d^5x \left(\frac{1}{4} \psi_{ab}^{(s)} \psi_{ij}^{(s)} \eta^{ai} \eta^{bj} - e^{-\phi} \sqrt{-\det(\gamma_{ab} + F_{ab})} \right)$$
 (3.37)
$$- \int (C_1 + C_3 + C_5') \wedge e^F,$$

where

$$F_{ab} = \partial_a A_b - \partial_b A_a + b_{ab}, \qquad (3.38)$$

$$C_5' = C_5 - \frac{1}{2}B \wedge C_3, \tag{3.39}$$

$$\psi_{ab}^{(s)} = \partial_a A_b^{(s)} - \partial_b A_a^{(s)}. \tag{3.40}$$

The action (3.37) describes the complete D4-brane [26] in type IIA supergravity background decoupled with unphysical field.

At this point, let us compare with the result from the PST formalism. It can be seen from the discussion in the literature that double dimensional reduction of the M5-brane action in the original PST action [4], [6], [7] on a circle is given by the dual D4-brane action, which can be realised as a duality transformation [26] of the D4-brane action. Alternatively, if one starts with the dual formulation of the M5-brane action [5],

[13], then it can be shown that double dimensional reduction on a circle directly gives rise to the D4-brane action. As for the case of Sen formalism, we have seen in this subsection that essentially by dualising some components of P, we obtain as a result of double dimensional reduction of the complete M5-brane action on a circle, the D4-brane action. Therefore, it seems that the original PST M5-brane action naturally gives rise, after double dimensional reduction on a circle, to dual D4-brane action. On the other hand, the Sen M5-brane action seems to naturally give rise to D4-brane action. In fact, as will be demonstrated in the next subsection, it can be seen that the Sen M5-brane action also naturally give rise to dual D4-brane action.

3.3 Dual D4-brane action from Sen M5-brane action

We have learned from the previous section that the key steps of double dimensional reduction of the M5-brane action in Sen formalism to the D4-brane action are essentially to dualise components P_{ab} of P and then eliminate Q. The five-dimensional DBI gauge field A and unphysical field $A^{(s)}$ then arise as linear combinations of the Lagrange multiplier and the remaining components P_{a5} of P.

In fact, by slightly modifying the procedure, it is possible to obtain the dual D4-brane action by dimensionally reducing Sen M5-brane action. The modification is simply to switch the role of the components of P so that we dualise components P_{a5} and as to be seen at the end of the process that components of P_{ab} along with the Lagrange multiplier will be linearly combined to form gauge field and unphysical field.

Let us show this in more details. We follow the same steps as those of the previous section up to eq.(3.21). Then, by defining

$$Y_{ab} = 2\partial_{[a}P_{b]5} \tag{3.41}$$

and introducing Lagrange multiplier term, we obtain

$$\tilde{S} = \frac{1}{2} \int d^5x \left(\frac{3}{4} \partial_a P_{bc} \partial^{[a} P^{bc]} + \frac{1}{4} Y_{ab} Y^{ab} + q^{ab} Y_{ab} \right)$$

$$+ \frac{1}{2} q_{ab} \partial_m P_{np} \epsilon^{mnpab} + \frac{1}{4} Y_{ab} \partial_m K_{np} \epsilon^{abmnp}$$

$$+ \frac{1}{12} \sqrt{-g} U + [q(R + \tilde{C})]$$

$$+ \frac{1}{2} [(q + R)(b - \tilde{C})] - \int C_5.$$
(3.42)

Integrating out Y gives

$$\widetilde{S} = \frac{1}{2} \int d^5x \left(\left[(\widetilde{\partial K} - \widetilde{\partial P} + q)^2 \right] + 2 \left[\widetilde{\partial K} \widetilde{\partial P} \right] \right.$$

$$\left. + \frac{1}{12} \sqrt{-g} U + \left[q(R + \widetilde{C}) \right] \right.$$

$$\left. + \frac{1}{2} \left[(q + R)(b - \widetilde{C}) \right] \right) - \int C_5,$$

$$(3.43)$$

where $\widetilde{\partial K}$ and $\widetilde{\partial P}$ are matrices with components $\widetilde{\partial K}^{mn} \equiv \epsilon^{mnijk} \partial_i K_{jk}/4$ and $\widetilde{\partial P}^{mn} \equiv \epsilon^{mnijk} \partial_i P_{jk}/4$, respectively. Indices for matrices $\widetilde{\partial K}, \widetilde{\partial P}$ are raised and lowered by the five-dimensional flat metric η . Let us define

$$\chi \equiv \widetilde{\partial P} - \widetilde{\partial K}, \qquad \chi^{(s)} \equiv \widetilde{\partial P} + \widetilde{\partial K}, \qquad (3.44)$$

which can be viewed as the field redefinitions

$$P_{ij} - K_{ij} \equiv 2B_{ij}, \qquad P_{ij} + K_{ij} \equiv 2B_{ij}^{(s)}.$$
 (3.45)

So χ and $\chi^{(s)}$ are dual of field strengths of B and $B^{(s)}$ with respect to five-dimensional flat metric. The action then becomes

$$\tilde{S} = \frac{1}{2} \int d^5x \left(\left[(-\chi + q)^2 \right] - \frac{1}{2} [\chi^2] + \frac{1}{2} [(\chi^{(s)})^2] + \frac{1}{12} \sqrt{-g} U + [q(R + \tilde{C})] + \frac{1}{2} [(q + R)(b - \tilde{C})]) - \int C_5.$$
(3.46)

Variation with respect to q gives

$$R = \chi - q. \tag{3.47}$$

Eliminating R from the action gives

$$\tilde{S} = \int d^5x \left(\frac{1}{4} [(\chi^{(s)})^2] + \frac{1}{4} [\chi^2] + \frac{1}{2} [q(-\chi + \tilde{C})] + \frac{1}{24} \sqrt{-g} U + \frac{1}{4} [\chi b] - \frac{1}{4} [\chi \tilde{C}] \right) - \int C_5.$$
(3.48)

The next step is to reexpress U. For this, let us define

$$\hat{F}^{a}{}_{b} \equiv \frac{F^{a}{}_{b}{}^{5}}{\sqrt{g^{55}}}, \qquad \hat{\tilde{F}}^{a}{}_{b} \equiv \frac{\tilde{F}^{a}{}_{b}{}^{5}}{\sqrt{g^{55}}}.$$
 (3.49)

Let indices of \hat{F}^{ab} and $\hat{\tilde{F}}^{ab}$ be lowered by γ_{ab} whereas indices of $F^{\mu\nu\rho}$ and $\tilde{F}^{\mu\nu\rho}$ are lowered by $g_{\mu\nu}$. We obtain

$$F_{\mu\nu\rho}F^{\mu\nu\rho} = -3[\hat{F}^2] + 3[\hat{\tilde{F}}^2].$$
 (3.50)

We need to eliminate \hat{F}_{ab} using nonlinear self-duality condition

$$\hat{F}^{a}{}_{b} = \frac{(1+z_{1})\hat{F}^{a}{}_{b} - (\hat{F}^{3})^{a}{}_{b}}{\sqrt{1+z_{1} + \frac{1}{2}z_{1}^{2} - z_{2}}},$$
(3.51)

where

$$z_1 \equiv \frac{1}{2}[\hat{\tilde{F}}^2], \qquad z_2 \equiv \frac{1}{4}[\hat{\tilde{F}}^4].$$
 (3.52)

This gives

$$U = -\frac{12(2+z_1)}{\sqrt{1+z_1 + \frac{1}{2}z_1^2 - z_2}}.$$
 (3.53)

Let us eliminate q from the action (3.46). By considering the ab5 component of H = Q - R, we obtain

$$q_{ab} - R_{ab} + b_{ab} = \frac{1}{\sqrt{g^{55}}} \left(g_{ac} \hat{F}^{c}{}_{b} - \frac{1}{2} \sqrt{-g} \epsilon_{abcmn} g^{c5} \hat{\tilde{F}}^{m}{}_{p} g^{pn} \right).$$
(3.54)

So along with eq.(3.47), we obtain

$$q_{ab} = \frac{1}{2} \left(\frac{g_{ac}}{\sqrt{g^{55}}} \hat{F}^c{}_b - \frac{1}{2} \frac{\sqrt{-g}}{\sqrt{g^{55}}} \epsilon_{abcmn} g^{c5} \hat{\tilde{F}}^m{}_p g^{pn} - b_{ab} + \sqrt{-g} \sqrt{g^{55}} \hat{\tilde{F}}^m{}_p \tilde{g}^{pn} \eta_{ma} \eta_{nb} + \tilde{C}_{ab} \right),$$
(3.55)

which completely determine Q in terms of other fields due to eq.(3.51). Here

$$\tilde{g}^{ab} \equiv g^{ab} - \frac{g^{a5}g^{b5}}{g^{55}}. (3.56)$$

We also have

$$\chi^{ab} = \sqrt{-g} \sqrt{g^{55}} \hat{\tilde{F}}^{a}{}_{c} \tilde{g}^{cb} + \tilde{C}^{ab}. \tag{3.57}$$

Let us define

$$W^{ab} \equiv \sqrt{-g} \sqrt{g^{55}} \hat{F}^a{}_a \tilde{g}^{db}$$
$$= -\frac{1}{3!} \epsilon^{abijk} F_{ijk}, \tag{3.58}$$

which is the dualisation of F_{ijk} with respect to five-dimensional flat metric η . The components of W are raised and lowered by η .

So the action becomes

$$\begin{split} \tilde{S} &= \int d^5x \left(\frac{3}{4} \partial_{[a} b^{(s)}_{bc]} \partial_{[i} b^{(s)}_{jk]} \eta^{ai} \eta^{bj} \eta^{ck} \right. \\ &- e^{-\phi} \sqrt{-\gamma} \sqrt{1 + z_1 + \frac{1}{2} z_1^2 - z_2} \\ &+ \frac{1}{2} [Wb] + \frac{1}{4} [\tilde{C}b] \\ &- \frac{1}{8} \frac{1}{g^{55}} \epsilon_{abcmn} g^{c5} W^{mn} W^{ab} \right) - \int C_5 \\ &= \int d^5x \left(\frac{3}{4} \partial_{[a} b^{(s)}_{bc]} \partial_{[i} b^{(s)}_{jk]} \eta^{ai} \eta^{bj} \eta^{ck} \right. \\ &- e^{-\phi} \sqrt{-G} \\ &- \frac{1}{24} \frac{\gamma^{cd} C_d \epsilon^{abmnp} F_{mnp} F_{abc}}{e^{-4\phi/3} + e^{2\phi/3} C_i \gamma^{ij} C_j} \right) \\ &+ \int (b \wedge (F - C_3) - C_5'), \end{split}$$
(3.59)

where

$$G_{ab} \equiv \gamma_{ab} - \frac{i}{6} e^{\phi} \frac{\gamma_{ac} \epsilon^{cdijk} F_{ijk} (\gamma_{db} + e^{2\phi} C_d C_b)}{\sqrt{-\gamma} \sqrt{1 + e^{2\phi} C_m \gamma^{mn} C_n}},$$
(3.60)

and, due to eq.(3.44), (3.57)-(3.58),

$$F_{ijk} = \partial_i B_{jk} + \partial_j B_{ki} + \partial_k B_{ij} + C_{ijk}. \quad (3.61)$$

The action (3.59) describes the complete dual D4-brane action [26] decoupled with unphysical field.

3.4 Duality from the viewpoint of Sen M5-brane action

It is well known that the D4-brane part of the action (3.37) and the dual D4-brane part of the action (3.59) are related by duality transformation. This can be shown [26] by considering the dualisation of A_a in the D4-brane action, which then gives rise to the dual D4-brane action. More explicitly, one replaces $2\partial_{[a}A_{b]}$ in the D4-brane action by ψ_{ab} , treats the latter as an independent field, introduces a 3-form Lagrange multiplier χ which puts $\psi_{ab} = 2\partial_{[a}A_{b]}$, then finally integrates out A_a and ψ_{ab} . This process gives rise to the dual D4-brane action, in which the Lagrange multiplier χ becomes the field strength of the 2-form field B.

It can be shown from the viewpoint of dimensionally reduced Sen M5-brane action that the action (3.37) and (3.59) are dual to each other. The prove can be performed quite simply by noting that the action (3.37) is equivalent to the action (3.24), whereas the action (3.59) is equivalent to the action (3.43). Let us replace $(\partial K)_{ab}$ and $(\partial P)_{ab}$ in the action (3.24) by $\Phi^{(1)}_{ab}$ and $\Phi^{(2)}_{ab}$, respectively. Let us treat $\Phi^{(1)}_{ab}$ and $\Phi^{(2)}_{ab}$ as independent fields and impose Lagrange multipliers $\Theta^{(1)}_{ab}$, $\Theta^{(2)}_{ab}$ which impose $\partial_{[a}\Phi^{(1)}_{bc]} = 0 = \partial_{[a}\Phi^{(2)}_{bc]}$. So the action (3.24) becomes

$$S = \int d^5x \left(-\frac{1}{2} \left[(\Phi^{(1)} + \Phi^{(2)} + q)^2 \right] + \left[\Phi^{(1)} \Phi^{(2)} \right] \right.$$
$$\left. + \frac{1}{4} \epsilon^{abcij} (\partial_a \Theta_{bc}^{(1)} \Phi_{ij}^{(1)} + \partial_a \Theta_{bc}^{(2)} \Phi_{ij}^{(2)}) \right.$$
$$\left. + \frac{1}{24} \sqrt{-g} U + \frac{1}{2} [q(R+b)] \right.$$
$$\left. - \frac{1}{4} [(q-R)(b-\tilde{C})] \right) - \int C_5.$$
(3.62)

Euler-Lagrange equations for $\Theta_{ab}^{(1)}$ and $\Theta_{ab}^{(2)}$ are $\partial_{[i}\Phi_{jk]}^{(1)}=0=\partial_{[i}\Phi_{jk]}^{(2)}$, which if the five-dimensional spacetime is topologically trivial in the sense that all closed forms are exact, then we can write $\Phi_{ij}^{(1)}=\partial_{[i}K_{j]},\Phi_{ij}^{(2)}=\partial_{[i}P_{j]},$ which upon substitution into eq.(3.62), we reobtain eq.(3.24) as expected. Let us however consider Euler-Lagrange equations for $\Phi_{ab}^{(1)}$ and $\Phi_{ab}^{(2)}$ which imply

$$(\Phi^{(1)})^{ab} = -q^{ab} - \frac{1}{4} \epsilon^{ijkab} \partial_i \Theta_{jk}^{(1)}, \qquad (3.63)$$

$$(\Phi^{(2)})^{ab} = -q^{ab} - \frac{1}{4} \epsilon^{ijkab} \partial_i \Theta_{jk}^{(2)}.$$
 (3.64)

Substituting these results into eq.(3.62), keeping in mind that indices for $q, \Phi^{(1)}, \Phi^{(2)}$ are raised and lowered by η , and denoting $\Theta_{ij}^{(1)} = -P_{ij}$ and $\Theta_{ij}^{(2)} = K_{ij}$, we obtain eq.(3.43). This finishes the proof in the viewpoint of Sen formalism that the D4-brane action and the dual D4-brane action are dual to each other.

4 Towards dimensional reduction of the Sen M5-brane action on other spaces

In [18], [20], analyses of dimensional reductions of the 6d Sen quadratic action are given. The examples of the spaces are a circle, a torus, K3, and a non-compact Riemann surface. Furthermore, in section 3 we have demonstrated that when non-linearising to the Sen M5-brane action, the dimensional reduction on a circle can naturally be given. In particular, it is natural to give, based

on the procedure of double dimensional reduction, either D4-brane action or dual D4-brane action. Note that these results suggest a remarkable feature of the Sen formalism that although its quadartic action couples to gravity in a complicated way, dimensional reduction on various cases of spaces can naturally be performed.

In contrast, in the PST formalism, although dimensional reduction can in principle be performed on any spaces, it is more natural to use specific action for dimensional reduction on a specific space. For example, it is natural to perform dimensional reduction of the original PST action [4], [6], [7] and its dual version [5], [13] on a circle giving rise respectively to D4-brane and dual D4-brane actions, whereas it is natural to perform dimensional reduction of the 3+3 M5-brane action [12] on T^3 to give an M2-brane action.

In this section, we will work towards generalising the analysis in section 3 so that it applies also to dimensional reductions on other spaces. We will only focus on special cases which contain some particular ways of dimensional reductions. We will also only give an outline of the analysis. The explicit details will be studied as future works. We expect that the analysis in this section could be served as a base for further generalisations, for example, double dimensional reduction of the complete M5-brane action on T^{5-p} to give Dp-brane with p < 4 (the case p = 4 has been given in the previous sections).

4.1 An outline of the analysis

Let us consider dimensional reduction of the Sen M5-brane action (2.7) from a 6 dimensional spacetime $\mathcal{M}_6 = \mathcal{D}_D \times \mathcal{C}_{6-D}$ with coordinates $x^{\mu}, \mu = 0, 1, 2, \cdots, 5$ to a D dimensional topologically trivial spacetime \mathcal{D}_D with coordinates $x^a, a = 0, 1, \cdots, D-1$. Let fields be independent from the coordinates $x^{\alpha}, \alpha = D, D+1, \cdots, 5$ of the compact space \mathcal{C}_{6-D} . Denote d and \hat{d} to be the exterior derivative on \mathcal{M}_6 and \mathcal{D}_D , respectively.

On \mathcal{M}_6 , let us denote a 3-form coordinate basis in six dimensions as $\mathcal{B} = \{dx^{\mu} \wedge dx^{\nu} \wedge dx^{\rho} \mid \mu, \nu, \rho \in \mathcal{B}\}$

 $^{^2 \}mathrm{In}$ this section, middle Greek alphabets for example μ, ν, ρ are used as indices of coordinates on \mathcal{M}_6 . Roman alphabets for example a,b,c,i,j,k,m,n are used as indices of coordinates on \mathcal{D}_D . Beginning Greek alphabets for example α,β are used as indices of coordinates on \mathcal{C}_{6-D} .

 $\{0,1,2,\cdots,5\}$. Let us pick 2n projection operators \mathcal{P}_I , $I=1,2,\cdots,2n$ such that

$$\mathcal{P}_{I}(dx^{\mu} \wedge dx^{\nu} \wedge dx^{\rho})$$

$$= \begin{cases} dx^{\mu} \wedge dx^{\nu} \wedge dx^{\rho} & \text{if } dx^{\mu} \wedge dx^{\nu} \wedge dx^{\rho} \in \mathcal{B}_{I}, \\ 0 & \text{if } dx^{\mu} \wedge dx^{\nu} \wedge dx^{\rho} \notin \mathcal{B}_{I}, \end{cases}$$

$$(4.1)$$

where $\mathcal{B}_1, \mathcal{B}_2, \dots, \mathcal{B}_{2n}$ are mutually exclusive subsets of $\mathcal{B} = \bigcup_{I=1}^{2n} \mathcal{B}_I$. Therefore, the projection operators satisfy

$$\sum_{I=1}^{2n} \mathcal{P}_I = 1, \tag{4.2}$$

$$*'\mathcal{P}_I *' = Y_{IJ}\mathcal{P}_J, \tag{4.3}$$

where Y_{IJ} are constants. Let us also impose a condition that the projection operators \mathcal{P}_I satisfy the property that for any 2-form A which is independent from coordinates of \mathcal{C}_{6-D} , there exists a 2-form A_I such that

$$\mathcal{P}_I dA = \hat{d}A_I. \tag{4.4}$$

Since A is independent from x^{α} , it can easily be seen that A_{I} are also independent from x^{α} . Let us require furthermore that the first n projection operators $\mathcal{P}_{\tilde{I}}, \hat{I} = 1, 2, \cdots, n$, are related to the last n projection operators $\mathcal{P}_{\tilde{I}}, \bar{I} = n + 1, n + 2, \cdots, 2n$, by

$$*'\mathcal{P}_{\hat{I}}*' = \delta_{\hat{I}-\bar{I}-n}\mathcal{P}_{\bar{I}}. \tag{4.5}$$

Therefore.

$$Y_{\bar{I}\bar{J}} = Y_{\hat{I}\hat{I}} = 0, \quad Y_{\hat{I}\bar{I}} = Y_{\bar{I}\hat{I}} = \delta_{\hat{I}\bar{I}-n}.$$
 (4.6)

Let us replace dP in eq.(2.7) by X and introduce Lagrange multiplier \tilde{P} which puts dX=0. The action becomes

$$S = \frac{1}{2} \int \left(\frac{1}{2} X \wedge *' X - 2Q \wedge X - d\tilde{P} \wedge X + \frac{d^6 x}{12} \sqrt{-g} U + Q \wedge R + F \wedge C_3 + 2C_6 \right).$$

$$(4.7)$$

Euler-Lagrange equation for X is given by

$$X + 2Q + *'d\tilde{P} = 0. (4.8)$$

As suggested by [27], if we substitute X from eq.(4.8) into eq.(4.7), we would obtain the action as in eq.(2.7) with P replaced by \tilde{P} . This is not the resulting action we are interested in. So let us instead only substitute the components $\mathcal{P}_{\tilde{J}}X$ from eq.(4.8):

$$\mathcal{P}_{\bar{I}}X = -2\mathcal{P}_{\bar{I}}Q - \mathcal{P}_{\bar{I}} *' d\tilde{P}. \tag{4.9}$$

into eq.(4.7). This gives

$$S = \frac{1}{2} \int \left(2\mathcal{P}_{\hat{I}} Q \wedge *' \mathcal{P}_{\hat{I}} Q + 2\mathcal{P}_{\hat{I}} d\tilde{P} \wedge *' \mathcal{P}_{\hat{I}} Q \right)$$

$$+ \frac{1}{2} \mathcal{P}_{\hat{I}} d\tilde{P} \wedge *' \mathcal{P}_{\hat{I}} d\tilde{P} + \frac{1}{2} \mathcal{P}_{\hat{I}} X \wedge *' \mathcal{P}_{\hat{I}} X$$

$$- 2 *' \mathcal{P}_{\hat{I}} Q \wedge \mathcal{P}_{\hat{I}} X - Y_{J\hat{I}} \mathcal{P}_{\bar{J}} d\tilde{P} \wedge \mathcal{P}_{\hat{I}} X$$

$$+ \frac{d^6 x}{12} \sqrt{-g} U + Q \wedge R + F \wedge C_3 + 2C_6 \right). \tag{4.10}$$

Note that if ω is a differential p-form on \mathcal{M}_6 and is independent on coordinates x^{α} of \mathcal{C}_{6-D} , we have $d\omega = \hat{d}\omega \equiv dx^a \wedge \partial_a \omega$. In particular, if $\hat{d}\omega = 0$, then by expressing in index-notation and using the fact that the D-dimensional spacetime is topologically trivial, it can be shown, for example by expressing in index notation, that there exists a (p-1)-form ϕ on \mathcal{M}_6 such that it is independent on x^{α} and that $\omega = \hat{d}\phi$.

The Euler-Lagrange equations for \tilde{P} are

$$\hat{d}(2\mathcal{P}_{\bar{I}}\mathcal{P}_{\bar{I}}Q + *'\mathcal{P}_{\hat{I}}\mathcal{P}_{\hat{I}}d\tilde{P} - \mathcal{P}_{\hat{I}}\mathcal{P}_{\hat{I}}X) = 0, \quad (4.11)$$

which imply

$$2\mathcal{P}_{\bar{i}}\mathcal{P}_{\bar{i}}Q + *'\mathcal{P}_{\hat{i}}\mathcal{P}_{\hat{i}}d\tilde{P} - \mathcal{P}_{\hat{i}}\mathcal{P}_{\hat{i}}X = -\hat{d}P, \quad (4.12)$$

where we denote the arbitrary x^{α} -independent 2—form on the 6-dimensional spacetime as P. Let us apply $\mathcal{P}_{\hat{I}}$ on eq.(4.12). This gives

$$\mathcal{P}_{\hat{I}}X = \mathcal{P}_{\hat{I}}\hat{d}P. \tag{4.13}$$

By substituting eq.(4.13) into eq.(4.10), we obtain

$$S = \frac{1}{2} \int \left(2\mathcal{P}_{\hat{I}} Q \wedge *' \mathcal{P}_{\hat{I}} Q - 2 *' \mathcal{P}_{\hat{I}} Q \wedge \mathcal{P}_{\hat{I}} (dP + d\tilde{P}) \right)$$

$$+ \frac{1}{2} \mathcal{P}_{\hat{I}} d\tilde{P} \wedge *' \mathcal{P}_{\hat{I}} d\tilde{P} + \frac{1}{2} \mathcal{P}_{\hat{I}} dP \wedge *' \mathcal{P}_{\hat{I}} dP \right)$$

$$+ \frac{d^6 x}{12} \sqrt{-g} U + Q \wedge R + F \wedge C_3 + 2C_6 ,$$

$$(4.14)$$

where we discard the integral of total derivative.

The next task is to eliminate Q by solving its Euler-Lagrange equations and substituting back into the action (4.14). Let us argue that this is very difficult, if at all possible, to approach this directly. In principle, the Euler-Lagrange equations of Q would not be sufficient to completely determine Q. One should also realise self-duality and non-linear self-duality conditions. In particular, one should solve eq.(2.14) to determine R in terms of Q, g, C_3 . The resulting expression of Q is expected to be in terms of P, P, Q, C_3 . It is likely that this expression would be complicated so that it is not even clear if it can be expressible in a closed form.

With the insight gained from the analysis in section 3, one may approach this strategically. Let us first consider the Euler-Lagrange equations for Q which can be expressed as

$$\mathcal{P}_{\hat{I}}R = \mathcal{P}_{\hat{I}}Q + \frac{1}{2}\mathcal{P}_{\hat{I}}(dP + d\tilde{P}). \tag{4.15}$$

Note that one can see from eq.(4.15) that some projections of $F = Q - R + C_3$ turn out to be expressible as simple expressions of P and \tilde{P} . That is

$$\mathcal{P}_{\hat{I}}F = -\frac{1}{2}\mathcal{P}_{\hat{I}}(dP + d\tilde{P}) + \mathcal{P}_{\hat{I}}C_3, \qquad (4.16)$$

which when using the property eq.(4.4), the first term on the RHS of eq.(4.16) is an exact differential form. One may use eq.(4.15) to express R in terms of Q then use eq.(2.14) to express Q in terms of $\mathcal{P}_{\hat{I}}F$. The idea is that the collection $\{\mathcal{P}_{\hat{I}}F\mid \hat{I}=1,2,\cdots,n\}$ contains ten components of F. The other ten components which are contained in the collection $\{\mathcal{P}_{\bar{I}}F\mid \bar{J}=n+1,n+2,\cdots,2n\}$ can in principle be expressed in terms of $\mathcal{P}_{\hat{I}}F$ by using non-linear self-duality relation of F. This means

that eq.(2.14) can be used to express

$$\mathcal{P}_{\bar{J}}F = \hat{V}_{\bar{J}}(\mathcal{P}_{\hat{I}}F, g), \tag{4.17}$$

where $\hat{V}_{\bar{J}}(\mathcal{P}_{\hat{I}}F, g)$ is a differential 3-form which is a function of $\mathcal{P}_{\hat{I}}F$ and g. Then by using $F = Q - R + C_3$ along with eq.(4.15)-(4.17), we obtain

$$\mathcal{P}_{\hat{I}}Q = \frac{1}{2}\mathcal{P}_{\hat{I}}F + \frac{1}{2}Y_{\hat{I}\bar{J}}*'\hat{V}_{\bar{J}}(\mathcal{P}_{\hat{I}}F,g) - \frac{1}{2}\mathcal{P}_{\hat{I}}(C_3 + *'C_3). \tag{4.18}$$

Let us then substitute eq.(4.15) and eq.(4.18) into the action (4.14). Note also that we may substitute the components $\mathcal{P}_{\bar{J}}F$ from eq.(4.17) into the expression of U(F,g) leaving only the dependency on $\mathcal{P}_{\hat{I}}F$ and g. More explicitly, we have after substitution,

$$U(F,g) = \hat{U}(\mathcal{P}_{\hat{I}}F,g). \tag{4.19}$$

The action (4.14) then becomes

$$S = \frac{1}{2} \int \left(\frac{1}{4} \mathcal{P}_{\hat{I}} (dP - d\tilde{P}) \wedge *' \mathcal{P}_{\hat{I}} (dP - d\tilde{P}) \right)$$

$$+ Y_{\hat{I}\bar{J}} \hat{V}_{\bar{J}} (\mathcal{P}_{\hat{I}} F, g) \wedge \mathcal{P}_{\hat{I}} F$$

$$+ \frac{d^6 x}{12} \sqrt{-g} \hat{U} (\mathcal{P}_{\hat{I}} F, g)$$

$$+ \mathcal{P}_{\hat{I}} \mathcal{P}_{\hat{I}} (2F - C_3) \wedge C_3 + 2C_6 \right).$$

$$(4.20)$$

Let us make field redefinitions

$$A = -\frac{1}{2}(P + \tilde{P}), \quad A^{(s)} = \frac{1}{2}(P - \tilde{P}).$$
 (4.21)

We may write

$$\mathcal{P}_{\hat{I}}dA = \hat{d}\hat{A}_{\hat{I}}, \quad \mathcal{P}_{\hat{I}}dA^{(s)} = \hat{d}\hat{A}_{\hat{I}}^{(s)}.$$
 (4.22)

Therefore,

$$S = \frac{1}{2} \int \left(\hat{d}\hat{A}_{\hat{I}}^{(s)} \wedge *' \hat{d}\hat{A}_{\hat{I}}^{(s)} + Y_{\hat{I}\bar{J}}\hat{V}_{\bar{J}}(\hat{F}_{\hat{K}}, g) \wedge \hat{F}_{\hat{I}} \right)$$

$$+ \frac{d^{6}x}{12} \sqrt{-g} \hat{U}(\hat{F}_{\hat{I}}, g)$$

$$+ \mathcal{P}_{\hat{I}}(2F_{\hat{I}} - \mathcal{P}_{\hat{I}}C_{3}) \wedge C_{3} + 2C_{6} ,$$

$$(4.23)$$

where $\hat{F}_{\hat{I}} \equiv \hat{d}\hat{A}_{\hat{I}} + \mathcal{P}_{\hat{I}}C_3$. After integrating out internal coordinates x^{α} the action (4.23) physical fields $\hat{A}_{\hat{I}}$ decoupled with free unphysical fields $\hat{A}_{\hat{I}}^{(s)}$. Generically, each of $\hat{A}_{\hat{I}}$ are collections of physical 0-form, 1-form, and 2-form fields in the D dimensional spacetime.

4.2 Duality

In subsection 3.4, we have seen that there is a simple way to realise the duality between D4-brane and dual D4-brane. From the point of view of the Sen M5-brane action, the duality is realised by simply switching the roles of components of P. In fact, it is natural to extend this idea to the cases of dimensional reduction to even lower dimensions.

Let us start by giving a quick summary to the result of subsection 4.1 as follows. Consider a dimensional reduction of the M5-brane action (2.7) to a D dimensional spacetime. The appropriate choice of 2n projection operators satisfy eq.(4.1)-(4.6) are also chosen. It turns out that only the first n projection operators $\mathcal{P}_1, \mathcal{P}_2, \dots, \mathcal{P}_n$ explicitly appear in the reduced action (4.23), which let us denote as $S_{1,2,3,\dots,n}$.

If we swap for example \mathcal{P}_1 with \mathcal{P}_{n+1} , the projection operators would still satisfy the properties (4.1)-(4.6). Therefore, we may obtain the reduced action $S_{n+1,2,3,\cdots,n}$ which is explicitly described by the projection operators $\mathcal{P}_{n+1}, \mathcal{P}_2, \mathcal{P}_3, \cdots, \mathcal{P}_n$.

In fact, the theories $S_{1,2,3,\dots,n}$ and $S_{n+1,2,3,\dots,n}$ are related by duality transformation. We first note that the action $S_{1,2,3,\dots,n}$ is equivalent to the action (4.14). Focusing on terms involving $\mathcal{P}_1Q, \mathcal{P}_1dP, \mathcal{P}_1d\tilde{P}$, we may express the action (4.14) as

$$S_{1,2,3,\cdots,n}$$

$$= \frac{1}{2} \int \left(2\mathcal{P}_1 Q \wedge *' \mathcal{P}_1 Q \right)$$

$$- 2 *' \mathcal{P}_1 Q \wedge \mathcal{P}_1 (dP + d\tilde{P})$$

$$+ \frac{1}{2} \mathcal{P}_1 d\tilde{P} \wedge *' \mathcal{P}_1 d\tilde{P}$$

$$+ \frac{1}{2} \mathcal{P}_1 dP \wedge *' \mathcal{P}_1 dP \right) + S',$$

$$(4.24)$$

where

$$S' = \frac{1}{2} \int \left(\sum_{\hat{I}=2}^{n} 2\mathcal{P}_{\hat{I}}Q \wedge *'\mathcal{P}_{\hat{I}}Q \right)$$

$$- \sum_{\hat{I}=2}^{n} 2 *'\mathcal{P}_{\hat{I}}Q \wedge \mathcal{P}_{\hat{I}}(dP + d\tilde{P})$$

$$+ \sum_{\hat{I}=2}^{n} \frac{1}{2}\mathcal{P}_{\hat{I}}d\tilde{P} \wedge *'\mathcal{P}_{\hat{I}}d\tilde{P}$$

$$+ \sum_{\hat{I}=2}^{n} \frac{1}{2}\mathcal{P}_{\hat{I}}dP \wedge *'\mathcal{P}_{\hat{I}}dP$$

$$+ \frac{d^{6}x}{12}\sqrt{-g}U + Q \wedge R + F \wedge C_{3} + 2C_{6} \right).$$

$$(4.25)$$

Note that by using the property (4.4) and the fact that P and \tilde{P} are independent from x^{α} , we may express

$$\mathcal{P}_1 dP = \hat{d}P_1, \qquad \mathcal{P}_1 d\tilde{P} = \hat{d}\tilde{P}_1. \tag{4.26}$$

In the action (4.24), let us set $\Phi \equiv \hat{d}P_1$, $\tilde{\Phi} \equiv \hat{d}\tilde{P}_1$ and introduce Lagrange multiplier to set $\hat{d}\Phi = \hat{d}\tilde{\Phi} = 0$ and $\mathcal{P}_1\Phi = \Phi$, $\mathcal{P}_1\tilde{\Phi} = \tilde{\Phi}$. So

$$S_{1,2,3,\cdots,n}$$

$$= \frac{1}{2} \int \left(2\mathcal{P}_1 Q \wedge *' \mathcal{P}_1 Q - 2 *' \mathcal{P}_1 Q \wedge (\Phi + \tilde{\Phi}) \right)$$

$$+ \frac{1}{2} \tilde{\Phi} \wedge *' \tilde{\Phi} + \frac{1}{2} \Phi \wedge *' \Phi$$

$$- \mathcal{P}_{n+1} dK \wedge \Phi - \mathcal{P}_{n+1} d\tilde{K} \wedge \tilde{\Phi} \right) + S',$$

$$(4.27)$$

where K and \tilde{K} are independent from x^{α} . Euler-Lagrange equations for Φ and $\tilde{\Phi}$ are

$$\Phi = -2\mathcal{P}_1 Q - *'\mathcal{P}_{n+1} dK, \quad \tilde{\Phi} = -2\mathcal{P}_1 Q - *'\mathcal{P}_{n+1} d\tilde{K}.$$
(4.28)

Substituting into the action (4.27) gives

$$S_{1,2,3,\dots,n}$$

$$= \frac{1}{2} \int \left(2\mathcal{P}_{n+1}Q \wedge *' \mathcal{P}_{n+1}Q \right)$$

$$- 2 *' \mathcal{P}_{n+1}Q \wedge \mathcal{P}_{n+1}(dK + d\tilde{K})$$

$$+ \frac{1}{2}\mathcal{P}_{n+1}d\tilde{K} \wedge *' \mathcal{P}_{n+1}d\tilde{K}$$

$$+ \frac{1}{2}\mathcal{P}_{n+1}dK \wedge *' \mathcal{P}_{n+1}dK \right) + S',$$

$$(4.29)$$

which after field redefinitions, it can easily be seen that

$$S_{1,2,3,\cdots,n} = S_{n+1,2,3,\cdots,n},$$
 (4.30)

as required.

The analysis above indeed generalises the analysis of section 3, in which D=5. So if we define $\mathcal{P}_1, \mathcal{P}_2$ such that

$$\mathcal{P}_{1}(dx^{\mu} \wedge dx^{\nu} \wedge dx^{\rho}) = 3\delta_{ij5}^{\mu\nu\rho} dx^{i} \wedge dx^{j} \wedge dx^{5},$$

$$\mathcal{P}_{2}(dx^{\mu} \wedge dx^{\nu} \wedge dx^{\rho}) = \delta_{ijk}^{\mu\nu\rho} dx^{i} \wedge dx^{j} \wedge dx^{k},$$
(4.31)

then the physical part of the action (4.23) describes a D4-brane. But if we define

$$\mathcal{P}_{1}(dx^{\mu} \wedge dx^{\nu} \wedge dx^{\rho}) = \delta^{\mu\nu\rho}_{ijk} dx^{i} \wedge dx^{j} \wedge dx^{k},$$

$$\mathcal{P}_{2}(dx^{\mu} \wedge dx^{\nu} \wedge dx^{\rho}) = 3\delta^{\mu\nu\rho}_{ij5} dx^{i} \wedge dx^{j} \wedge dx^{5},$$
(4.32)

then the physical part of the action (4.23) describes a dual D4-brane.

By similar considerations, any pair $(\mathcal{P}_k, \mathcal{P}_{k+n}), k = 1, 2, \dots, n$ of projection operators can be swapped and that it is also possible to swap more than one pair at a time. The resulting action is dual to the original action. For simplicity, let us discuss an example case where the action is quadratic. In this case, the action (4.23) is

$$S = \frac{1}{2} \int \left(\hat{d}\hat{A}_{\hat{I}}^{(s)} \wedge *' \hat{d}\hat{A}_{\hat{I}}^{(s)} + Y_{\hat{I}\bar{J}}\hat{V}_{\bar{J}}(\hat{F}_{\hat{K}}, g) \wedge \hat{F}_{\hat{I}} \right) + \mathcal{P}_{\hat{I}}(2\hat{F}_{\hat{I}} - \mathcal{P}_{\hat{I}}C_3) \wedge C_3 + 2C_6,$$

$$(4.33)$$

where the form of $\hat{V}_{\bar{J}}(\hat{F}_{\hat{I}},g)$ will be given on a case by case basis by rearranging the 6d *-self-duality condition of F.

In particular, let us consider an example case in which D=4, the compact space is a torus, and the 6d metric is $g=g_4 \oplus g_2$. Let us pick the projection operators as follows

$$\mathcal{P}_{1}(dx^{\mu}\wedge dx^{\nu}\wedge dx^{\rho}) = 6\delta_{i45}^{\mu\nu\rho}dx^{i}\wedge dx^{4}\wedge dx^{5},$$

$$\mathcal{P}_{2}(dx^{\mu}\wedge dx^{\nu}\wedge dx^{\rho}) = 3\delta_{ij4}^{\mu\nu\rho}dx^{i}\wedge dx^{j}\wedge dx^{4},$$

$$\mathcal{P}_{3}(dx^{\mu}\wedge dx^{\nu}\wedge dx^{\rho}) = \delta_{ijk}^{\mu\nu\rho}dx^{i}\wedge dx^{j}\wedge dx^{k},$$

$$\mathcal{P}_{4}(dx^{\mu}\wedge dx^{\nu}\wedge dx^{\rho}) = 3\delta_{ij5}^{\mu\nu\rho}dx^{i}\wedge dx^{j}\wedge dx^{5}.$$

$$(4.34)$$

Furthermore, let us consider a consistent truncation by setting

$$\hat{F}_1 = \hat{F}_3 = \mathcal{P}_1 C_3 = \mathcal{P}_3 C_3 = 0, \tag{4.35}$$

or in component form

$$F_{i45} = F_{ijk} = C_{i45} = C_{ijk} = 0.$$
 (4.36)

This example case is exactly what studied before in [18] in which $g_4 = \eta_4$ is the flat 4d metric. Here, however, we shall generalise this by taking g_4 to be a general 4d metric which depends on the 4d coordinates x^i . The *-self-duality condition of F in 6d gives

$$F^{ij4} = \frac{1}{g_{44}} \left(g^{mi} g^{nj} + g_{45} \frac{1}{2\sqrt{-g}} \epsilon^{ijmn} \right) F_{mn4},$$
(4.37)

where $F_{mn4} = 2\partial_{[m}A_{n]4} + C_{mn4}$. Therefore, the physical part of the action (4.33) is

$$S_{phys} = \int d^4x \left(-\frac{1}{4} \frac{\sqrt{g_{44}g_{55} - g_{45}^2}}{g_{44}} \sqrt{-g_4} F_{ij4} F_{mn4} g^{im} g^{jn} - \frac{1}{8} \frac{g_{45}}{g_{44}} \epsilon^{mnij} F_{ij4} F_{mn4} \right) + \int d^4x \left(-\frac{1}{4} F_{ij4} C_{mn5} \epsilon^{ijmn} - \frac{1}{8} C_{ij4} C_{mn5} \epsilon^{ijmn} - \frac{1}{4!} C_{ijmn45} \epsilon^{ijmn} \right),$$

$$(4.38)$$

where ϵ^{ijmn} is a 4d Levi-Civita tensor which is defined as $\epsilon^{ijmn} = \epsilon^{ijmn45}$, and for simplicity, we set $\int dx^4 = \int dx^5 = 1$. By making the swap $\mathcal{P}_2 \leftrightarrow$

 \mathcal{P}_4 , we obtain the dual action

$$\tilde{S}_{phys} = \int d^4x \left(-\frac{1}{4} \frac{\sqrt{g_{44}g_{55} - g_{45}^2}}{g_{55}} \sqrt{-g_4} F_{ij5} F_{mn5} g^{im} g^{jn} + \frac{1}{8} \frac{g_{45}}{g_{55}} \epsilon^{mnij} F_{ij5} F_{mn5} \right) + \int d^4x \left(\frac{1}{4} F_{ij5} C_{mn4} \epsilon^{ijmn} + \frac{1}{8} C_{ij4} C_{mn5} \epsilon^{ijmn} - \frac{1}{4!} C_{ijmn45} \epsilon^{ijmn} \right).$$
(4.39)

The action (4.38) and its dual (4.39) transform to each other under the S-duality transformation

$$g_{44} \mapsto g_{55}, \quad g_{45} \mapsto -g_{45}, \quad g_{55} \mapsto g_{44}, \quad (4.40)$$

$$F_{ij4} \mapsto -F_{ij5}, \quad F_{ij5} \mapsto F_{ij4},$$

$$C_{ij4} \mapsto -C_{ij5}, \quad C_{ij5} \mapsto C_{ij4}.$$

$$(4.41)$$

Note that the transformation (4.40) is equivalent to

$$\tau \mapsto -\frac{1}{\tau},\tag{4.42}$$

where

$$\tau = \frac{-g_{45} + i\sqrt{g_{44}g_{55} - g_{45}^2}}{g_{44}}. (4.43)$$

The transformation (4.42) is what expected from the conformal symmetry of the 6d action of self-dual 3-form [25]. Furthermore, in the context of D3-brane, τ encodes axion and dilation as $\tau = C_0 + ie^{-\phi}$, whereas C_{ij4} and C_{ij5} are identified with B_2 and C_2 . The transformations (4.40)-(4.41) is precisely the S-duality transformation for D3-brane

It is clear that the reduced theories (4.38)-(4.39) we have obtained is expressed entirely in terms of standard fields. This is possible largely due to that we have chosen appropriate projection operators, in particular the ones which project to ij4 or ij5 components. On the other hand, for the choice studied in [18], in which the components $\hat{i}\hat{j}\alpha$ where $\hat{i},\hat{j}\in\{1,2,3\},\alpha\in\{4,5\}$ are chosen, it is not yet clear whether the Lagrangian can be expressed entirely in terms of standard fields.

Note that due to the expected form of the reduced theory under dimensional reduction and

how it transforms under S-duality, it could be expected that if one considers double dimensional reduction of the complete Sen M5-brane action (2.7) on a torus, then one could obtain the complete D3-brane action on a circle. Indeed, explicit checks are required. If this succeeds, one may then compare and contrast the approach with its counterpart [28] given in the PST formalism.

It is remarkable that the duality can easily be obtained simply by appropriately swapping the projection operators. In fact, since the analysis only involves mainly the field P and projection operators, similar analysis for the duality of dimensional reduction of self-dual (2p+1)-forms in 4p+2 dimensions in Sen formalism can easily be performed. This way that dualities of dimensionally reduced theories are realised in the Sen formalism is another property which makes the Sen formalism appealing. Of course, further efforts are still required to eliminate Q from each action to arrive at the final form.

5 Conclusion and Discussion

In this work, we have studied double dimensional reduction of the Sen M5-brane action (2.7). In particular, we have explicitly shown that double dimensional reduction of the Sen M5-brane action on a circle indeed gives, depending on how the projection operators are chosen, the complete D4-brane action or the complete dual D4-brane action. The procedure we have followed is essentially just a slight modification of that given in [20]. It is remarkable that although the Sen M5brane action couples to gravity in a complicated way and that there are fields which do not transform in the standard way under diffeomorphism, it seems that dimensional reduction can be carried out relatively simply giving rise to reduced actions which describe dynamics of fields which transform in a standard way under diffeormorphism.

We have also outlined an extension of dimensional reduction on other spaces. The analysis suggests that provided that one picks projection operators which satisfy eq.(4.1)-(4.6), the dimensional reduction of the Sen M5-brane action gives rise to the reduced action (4.23) which contains standard fields.

Note that one aspect which makes the procedure of dimensional reduction simple despite

the aforementioned complications of the Sen M5-brane action is that it allows one to easily integrating out Q. In particular, it avoids the need to explicitly express R in terms of Q,g,C_3 . As a future work, it would be interesting to see whether it is possible to extend and modify the procedure to show explicitly that the Sen M5-brane action is equivalent to the M5-brane actions in PST formalism. We anticipate that if this is at all possible, the procedure should also allow one to integrate out Q in a simple way.

We have found another remarkable feature of Sen M5-brane action that dualities of dimensional reduced action seem to be encoded within the Sen M5-brane action. For example, as demonstrated explicitly in this paper, in the viewpoint of string theory, it would take some efforts to show that the D4-brane and dual D4-brane actions are dual to each other. On the other hand, from the viewpoint of the Sen M5-brane action, the D4-brane and dual D4-brane actions can be put in a similar form. This can be seen by comparing eq.(3.24) and eq.(3.43), which can be transformed to each other simply by swapping the projection operators between eq.(4.31) and eq.(4.32). We have also shown that this swapping is a result of dualisation.

We have outlined a generalisation to dimensional reduction on other manifolds. As long as the projection operators can be given to satisfy (4.1)-(4.6), the process will give rise to the reduced action (4.23) whose physical part describe fields which transform under standard diffeomorphism. Apart from the D4-brane and dual D4-brane actions, it can be expected that the action (4.23) should also give rise to other complete actions in string theory and M-theory for example D3-brane, D2-brane, D1-brane, F1-brane, M2-brane actions and whether S-dualities in these cases are the results of simple swapings of the projection operators. For completeness, explicit checks should be made in each case.

The analysis for dimensional reduction on other spaces as given in section 4 heavily relies on the assumption that the projection operators should satisfy eq.(4.1)-(4.6). It would be interesting to see whether the analysis can be extended to the cases where some of the properties (4.1)-(4.6) are generalised. In any case, it would also be interesting to consider dimensional reduction of the Sen M5-brane action on other spaces. In particular, one might consider the dimensional reduction

of the Sen M5-brane action on Riemann surface and K3. This is to see non-linear extension to the result of [20].

It could also be expected that other kinds of dimensional reduction for the Sen M5-brane action could also easily be carried out. In particular, the null dimensional reduction on a (2,0) theory in the Sen formalism has been carried out in [21]. So one might try to use a similar method to study the null reduction of the Sen M5-brane action.

Although the analysis in section 4 is given explicitly for the six-dimensional Sen M5-brane action, the extension especially for the quadratic version which describes self-dual (2p+1)-form on 4p+2 dimensional spacetime can easily be done. In particular, for the case p=2, the Sen quadratic action describes type IIB supergravity. So it would also be interesting to study how the dimensional reduction of type IIB supergravity is realised in Sen formalism.

Acknowledgements

We would like to thank Sheng-Lan Ko for discussions. A.P. is supported by a Development and Promotion of Science and Technology Talents Project (DPST) scholarship from the Royal Thai Government. This version of the article has been accepted for publication, after peer review but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: http://dx.doi.org/10.1140/epjc/s10052-023-11892-2

Data Availability Statement No data was analysed or generated in this work.

References

- [1] Henneaux, M., Teitelboim, C.: Dynamics of Chiral (Selfdual) P Forms. Phys. Lett.
 B 206, 650-654 (1988) https://doi.org/10.1016/0370-2693(88)90712-5
- [2] Perry, M., Schwarz, J.H.: Interacting chiral gauge fields in six-dimensions and Born-Infeld theory. Nucl. Phys. B **489**,

- 47-64 (1997) https://doi.org/ 10.1016/S0550-3213(97)00040-0 arXiv:hep-th/ 9611065
- [3] Schwarz, J.H.: Coupling a selfdual tensor to gravity in six-dimensions. Phys. Lett. B **395**, 191–195 (1997) https://doi.org/10.1016/S0370-2693(97)00094-4 arXiv:hep-th/9701008
- [4] Aganagic, M., Park, J., Popescu, C., Schwarz, J.H.: World volume action of the M theory five-brane. Nucl. Phys. B 496, 191–214 (1997) https://doi.org/10.1016/S0550-3213(97)00227-7 arXiv:hep-th/9701166
- [5] Ko, S.-L., Vanichchapongjaroen, P.: The Dual Formulation of M5-brane Action. JHEP
 06, 022 (2016) https://doi.org/10.1007/ JHEP06(2016)022 arXiv:1605.04705 [hep-th]
- [6] Pasti, P., Sorokin, D.P., Tonin, M.: Covariant action for a D = 11 five-brane with the chiral field. Phys. Lett. B 398, 41–46 (1997) https://doi.org/10.1016/S0370-2693(97)00188-3 arXiv:hep-th/9701037
- [7] Bandos, I.A., Lechner, K., Nurmagambetov, A., Pasti, P., Sorokin, D.P., Tonin, M.: Covariant action for the superfivebrane of M theory. Phys. Rev. Lett. 78, 4332–4334 (1997) https://doi.org/10.1103/PhysRevLett.78.4332 arXiv:hep-th/9701149
- [8] Maznytsia, A., Preitschopf, C.R., Sorokin, D.P.: Duality of selfdual actions. Nucl. Phys. B 539, 438–452 (1999) https://doi.org/10.1016/S0550-3213(98)00741-X arXiv:hep-th/9805110
- [9] Maznytsia, A., Preitschopf, C.R., Sorokin, D.P.: Dual actions for chiral bosons. In: 10th Summer School / Seminar (Volga-10) on Recent Problems in Theoretical and Mathematical Physics (1998)
- [10] Chen, W.-M., Ho, P.-M.: Lagrangian Formulations of Self-dual Gauge Theories in Diverse Dimensions. Nucl. Phys. B 837, 1–21 (2010) https://doi.org/10.1016/j.nuclphysb. 2010.04.015 arXiv:1001.3608 [hep-th]
- [11] Huang, W.-H.: Lagrangian of Self-dual

- Gauge Fields in Various Formulations. Nucl. Phys. B **861**, 403–423 (2012) https://doi.org/10.1016/j.nuclphysb.2012.03.017 arXiv:1111.5118 [hep-th]
- [12] Ko, S.-L., Sorokin, D., Vanichchapongjaroen,
 P.: The M5-brane action revisited. JHEP
 11, 072 (2013) https://doi.org/10.1007/
 JHEP11(2013)072 arXiv:1308.2231 [hep-th]
- [13] Ko, S.-L., Vanichchapongjaroen, P.: A covariantisation of M5-brane action in dual formulation. JHEP 01, 072 (2018) https://doi.org/10.1007/JHEP01(2018)072 arXiv:1712.06408 [hep-th]
- [14] Mkrtchyan, K.: On Covariant Actions for Chiral p-Forms. JHEP **12**, 076 (2019) https://doi.org/10.1007/JHEP12(2019)076 arXiv:1908.01789 [hep-th]
- [15] Avetisyan, Z., Evnin, O., Mkrtchyan, K.: Nonlinear (chiral) p-form electrodynamics. JHEP 08, 112 (2022) https://doi.org/10.1007/JHEP08(2022)112 arXiv:2205.02522 [hep-th]
- [16] Evnin, O., Mkrtchyan, K.: Three approaches to chiral form interactions. Differ. Geom. Appl. 89, 102016 (2023) https://doi.org/10. 1016/j.difgeo.2023.102016 arXiv:2207.01767 [hep-th]
- [17] Sen, A.: Covariant Action for Type IIB Supergravity. JHEP 07, 017 (2016) https://doi.org/10.1007/JHEP07(2016)017 arXiv:1511.08220 [hep-th]
- [18] Sen, A.: Self-dual forms: Action, Hamiltonian and Compactification. J. Phys. A 53(8), 084002 (2020) https://doi.org/10.1088/1751-8121/ab5423 arXiv:1903.12196 [hep-th]
- [19] Lambert, N.: (2,0) Lagrangian Structures. Phys. Lett. B 798, 134948 (2019) https://doi.org/10.1016/j.physletb.2019.134948 arXiv:1908.10752 [hep-th]
- [20] Andriolo, E., Lambert, N., Papageorgakis, C.: Geometrical Aspects of An Abelian (2,0) Action. JHEP 04, 200 (2020) https://doi.org/10.1007/JHEP04(2020)200

arXiv:2003.10567 [hep-th]

- [21] Lambert, N., Orchard, T.: Null reductions of the M5-brane. JHEP 12, 037 (2020) https://doi.org/10.1007/JHEP12(2020)037 arXiv:2005.14331 [hep-th]
- [22] Gustavsson, A.: A nonabelian M5 brane Lagrangian in a supergravity background. JHEP 10, 001 (2020) https://doi.org/ 10.1007/JHEP10(2020)001 arXiv:2006.07557 [hep-th]
- [23] Vanichchapongjaroen, P.: Covariant M5-brane action with self-dual 3-form. JHEP
 05, 039 (2021) https://doi.org/10.1007/ JHEP05(2021)039 arXiv:2011.14384 [hep-th]
- [24] Andriolo, E., Lambert, N., Orchard, T., Papageorgakis, C.: A path integral for the chiral-form partition function. JHEP 04, 115 (2022) https://doi.org/10.1007/ JHEP04(2022)115 arXiv:2112.00040 [hep-th]
- [25] Witten, E.: Geometric Langlands From Six Dimensions (2009) arXiv:0905.2720 [hep-th]
- [26] Aganagic, M., Park, J., Popescu, C., Schwarz, J.H.: Dual D-brane actions. Nucl. Phys. B 496, 215–230 (1997) https:// doi.org/10.1016/S0550-3213(97)00257-5 arXiv:hep-th/9702133
- [27] Lambert, N.: Duality and fluxes in the sen formulation of self-dual fields. Phys. Lett. B
 840, 137888 (2023) https://doi.org/10.1016/j.physletb.2023.137888 arXiv:2302.10955 [hep-th]
- [28] Berman, D.: M5 on a torus and the three-brane. Nucl. Phys. B **533**, 317–332 (1998) https://doi.org/10.1016/S0550-3213(98)80009-6 arXiv:hep-th/9804115